## Table 198. Selected Respiratory Diseases Among Persons 18 Years of Age and Over by Selected Characteristics: 2009

[In thousands (227,371 represents 227,371,000). Respondents were asked in two separate questions if they had ever been told by a doctor or other health professional that they had emphysema or asthma. Respondents who had been told they had asthma were asked if they still had asthma. Respondents were asked in three separate questions if they had been told by a doctor or other health professional in the past 12 months that they had hay fever, sinusitis, or bronchitis. Based on the National Health Interview Survey, a sample survey of the civilian noninstitutionalized population; see Appendix III]

		Selected respiratory conditions <sup>1</sup>						
Selected characteristic	Total	Emphy-	Asthma		Hay		Chronic	
	persons	sema	Ever	Still	fever	Sinusitis	bronchitis	
Total <sup>2</sup>	227,371	4,895	29,734	17,456	17,738	29,305	9,908	
SEX								
Male Female	109,844 117,527	2,578 2,317	12,311 17,423	6,058 11,398	7,546 10,192	10,555 18,750	3,189 6,718	
AGE		1						
18 to 44 years 45 to 64 years 65 to 74 years 75 years and over	110,337 79,195 20,597 17,242	369 2,065 1,194 1,268	15,743 9,706 2,466 1,819	8,368 6,182 1,720 1,186	7,106 7,932 1,629 1,072	11,342 12,792 3,028 2,142	3,093 4,411 1,329 1,074	
RACE								
Race alone <sup>3</sup>	224,290 183,739 27,374	4,825 4,357 360	29,056 23,880 3,854	17,061 13,990 2,380	17,406 15,061 1,551	28,819 24,447 3,513	9,750 8,386 1,082	
American Indian or Alaska Native	1,856 10,763 558	(B) <sup>4</sup> 59 (B) <sup>4</sup> 70	275 968 (B)	4 177 490 4 25	4 104 670 (B)	155 694 (B)	<sup>4</sup> 78 184 (B)	
Two or more races 5	3,082	⁴ <b>`</b> 7Ó	678	394	332	486	158	
Hispanic or Latino	31,312 19,687	200 107	3,225 1,557	1,721 826	1,648 980	2,584 1,561	820 362	
Not Hispanic or Latino	196,059 155,185 26,213	4,696 4,173 350	26,509 21,035 3,675	15,735 12,484 2,281	16,090 13,532 1,497	26,721 22,123 3,393	9,088 7,647 1,046	

B Figure too small to meet statistical standards for reliability of a derived figure. ¹ A person may be represented in more than one column. ² Total includes other races not shown separately. ³ Refers to persons who indicated only a single race group. ⁴ Figure does not meet standard of reliability or precision. ⁵ Refers to all persons who indicated more than one race group. ⁶ Persons of Hispanic or Latino origin may be any race or combination of races.

Source: National Center for Health Statistics, Summary Health Statistics for the U.S. Population: National Health Interview Survey, 2009, Vital and Health Statistics, Series 10, Number 249, 2010. See also <a href="http://www.cdc.gov/nchs/data/series/sr\_10/sr10\_249.pdf">http://www.cdc.gov/nchs/data/series/sr\_10/sr10\_249.pdf</a>.

## Table 199. Persons 18 Years of Age and Over With Migraines and Pains in the Neck, Lower Back, Face, or Jaw by Selected Characteristics: 2009

[In thousands (227,371 represents 227,371,000). Based on National Health Interview Survey, a sample survey of the civilian noninstitutionalized population, Appendix III]

		Migraine and pain 1					
Selected characteristic		Migraine or		Pain in	Pain in		
Selected Characteristic	Total	severe	Pain in	lower	face or		
	persons	headache 2	neck 3	back 3	jaw 3		
Total 4	227,371	35,973	34,954	64,810	11,501		
SEX							
Male	109,844	11,098	14,095	28,842	3,654		
Female	117,527	24,875	20,859	35,968	7,847		
AGE							
18 to 44 years	110,337	21,706	14,306	26,973	5,752		
45 to 64 years	79,195	11,893	15,147	25,795	4,400		
65 to 74 years	20,597	1,415	3,136	6,198	799		
75 years and over	17,242	959	2,365	5,843	550		
RACE							
Race alone 5	224,290	35,295	34,355	63,689	11,213		
White	183,739	29,118	29,430	53,817	9,599		
Black or African American	27,374	4,811	3,569	7,286	1,115		
American Indian or Alaska Native	1,856	429	363	566	171		
Asian	10,763	902 6 33	935 <sup>6</sup> 57	1,879 <sup>6</sup> 141	303		
Two or more races 7	558 3,082	678	598	1,121	(B) 288		
	3,002	070	330	1,121	200		
HISPANIC ORIGIN AND RACE 8	04.040	F 000	4.044	7.005	4.045		
Hispanic or Latino	31,312 19.687	5,362 3.297	4,341 2.446	7,905 4.319	1,345 818		
Not Hispanic or Latino	196,059	3,297 30,611	30,613	4,319 56.905	10,156		
White, single race	155,185	24.379	25,497	46,665	8,423		
Black or African American, single race	26,213	4.546	3.426	6.982	1,065		

B Figure to small to meet statistical standards for reliability of a derived figure. ¹ A person may be represented in more than one column. ² Respondents were asked, "During the past 3 months, did you have a severe headache or migraine?" Respondents were instructed to report pain that had lasted a whole day or more and, conversely, not to report fleeting or minor aches or pains. ³ Respondents were asked, "During the past 3 months, did you have a neck pain; or low back pain; or facial ache or pain in the jaw muscles or the joint in front of the ear?" Respondents were instructed to report pain that had lasted a whole day or more and, conversely, not to report fleeting or minor aches or pains. ⁴ Total includes other races not shown separately. ⁵ Refers to persons who indicated only a single race group. ⁶ Figure does not meet standard of reliability or precision. ⁿ Refers to all persons who indicated more than one race group. ⁶ Persons of Hispanic or Latino origin may be any race or combination of races.

Source: National Center for Health Statistics, Summary Health Statistics for the U.S. Adults: National Health Interview Survey, 2009, Vital and Health Statistics, Series 10, Number 249, 2010. See also <a href="https://www.cdc.gov/nchs/data/series/sr\_10/sr10/sr10\_249.pdf">https://www.cdc.gov/nchs/data/series/sr\_10/sr10\_249.pdf</a>.